



## **Molasses injection as a MEOR strategy: Enrichment incubations of brine/oil from North Sea Oil Field**

**Halim, Amalia Yunita; Pedersen, Dorthe Skou; Eliasson Lantz, Anna; Nielsen, Sidsel Marie; Shapiro, Alexander**

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Halim, A. Y., Pedersen, D. S., Eliasson Lantz, A., Nielsen, S. M., & Shapiro, A. (2013). *Molasses injection as a MEOR strategy: Enrichment incubations of brine/oil from North Sea Oil Field*. Poster session presented at 4th International Symposium on Applied Microbiology and Molecular Biology in Oil Systems, Rio de Janeiro, Brazil.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Molasses injection as a MEOR strategy:

## Enrichment incubations of brine/oil from North Sea Oil Field


Amalia Halim\*, Dorthe Skou Pedersen\*\*, Anna Eliasson Lantz\*\*\*, Sidsel Marie Nielsen\*, Alexander Shapiro\*

\* Center for Energy Resources Engineering (CERE), Department of Chemical and Biochemical Engineering, DTU

\*\* Danish Technological Institute (DTI)

\*\*\* Centre for Microbial Biotechnology (CMB), Department of System Biology, DTU

### Background and Objectives



### Experimental Design






Figure 1. Anaerobic enrichment tests by using brine and crude oil from the North Sea oil field for 28 days at 55°C. Three different conditions on the enrichment test: (a) incubation A (control) – no nutrient addition, (b) incubation B with addition of 2% molasses and (c) incubation C with addition of 2% molasses and 2.5 mM nitrate.

#### Cell hydrophobicity



#### How many microbes ? qPCR

#### Which microbes? pyrosequencing



- Sugar hydrolysis and metabolite production
- Oil characteristics (viscosity, density, and total weight fraction)
- Interfacial tension (oil vs. brine)
- Emulsion



### Acknowledgement

- This study has been carried as a part of the BioRec project supported by the Danish National Advanced Technology Foundation, Mærsk Oil and Gas AS and DONG AS.
- Nordic Sugar is acknowledged for providing the molasses for the experiment.

### Contact Information

- Amalia Halim  
[amah@kt.dtu.dk](mailto:amah@kt.dtu.dk)  
Technical University of Denmark (DTU)  
Søtofts Plads, b 229, DK-2800 Lyngby
- Dorthe Skou Pedersen  
[dspe@teknologisk.dk](mailto:dspe@teknologisk.dk)  
DTI Oil and Gas  
Kongsvang Allé 29, DK-8000 Aarhus C

### Results




Figure 2. (a) molasses (sucrose) hydrolyses by microbes in incubation B (blue) and incubation C (red). Nitrate ( $\text{NO}_3^-$ ) was monitored in incubation C,  $\text{NO}_3^-$  was added at day 2, 6, 9, 14, 19 (orange circle) to keep  $\text{NO}_3^-$  concentration at 212.5 mg/l (2.5mM), (b) acetate production and brine pH during incubation, (c) ethanol production during incubation.

- Sucrose hydrolyses (incubations B and C) – presence of fermentative microbes. Nitrate was consumed continuously in incubation C (Fig. 2a).
- Acetate production – pH decrease (incubations B and C, Fig. 2b) and ethanol production (Fig. 2c).
- No production of succinate, lactate and glycerol on all incubations.




Figure 3. Microbes were present in the oil-brine interphase in all incubations (red arrow). (a) In incubation A, without molasses, microbes had a tendency to form aggregates. (b, c) In incubation B and C, with addition of molasses, microbes are present as single cell (loose particle).

Table 3. Interfacial tension (IFT) between the brine and the crude oil in incubation A, B and C

Sample	T0	T28 - with biomass	T 28 w/o biomass
A	19.29 ± 0.12	17.70 ± 1.48	15.20 ± 0.32
B	18.79 ± 0.18	2.34 ± 0.15	21.65 ± 0.15
C	18.60 ± 0.09	2.96 ± 0.43	22.60 ± 0.88

- In chalk it is important that microbes are present as single cells (hydrophilic) to prevent blocking of pore throat.
- No significant change in oil characteristics: viscosity, density and composition (total weight fraction) in all incubations.
- Formation of gasses, emulsions and smaller oil droplets in incubations B and C.
- IFT decreases when microbes are present in oil-brine interphase (Table 3).




Figure 4. Abundance of 16S rRNA bacteria and 16S rRNA archaea gene copies in incubations A, B and C.





Figure 5. Development in (a) bacterial community at phylum level and (b) archaeal community at organism level in incubation A, B and C.

Table 2. Summary of heat map of the most abundant microorganisms based on sequencing of the 16S rRNA gene of (a) bacteria and (b) archaea



- Microbial growth was stimulated by addition of molasses in incubations B and C (Fig. 4).
- Sequencing revealed the population dynamics of the microbial community (Fig. 5 and Table 2).

### Conclusion

- Production of fermentation products : acetate, ethanol and gas from sugar hydrolyses.
- No significant change in oil characteristics; but formation of emulsion, smaller oil droplets and reduction of IFT were observed due to microbial biomass when molasses was added (incubation B and C).
- Addition of molasses made microbe cells hydrophilic - prevent microbes to form aggregates.
- Molasses stimulated microbial growth in incubations B and C, as shown by increase in cell number of both bacteria and archaea (qPCR results).
- Pyrosequencing revealed that molasses stimulated the growth of *Petrotoga* and *Anaerobaculum*.
- Nitrate was consumed continuously and *Petrotoga* was favoured over *Anaerobaculum* in the presence of nitrate (incubation C).
- The growth of methanogenic archaea was potentially fueled by products from fermenting bacteria.